

5.308 ordered_global_cardinality

	DESCRIPTION	LINKS	GRAPH
Origin	[312]		
Constraint	<code>ordered_global_cardinality(VARIABLES, VALUES)</code>		
Usual name	<code>ordgcc</code>		
Synonym	<code>ordered_gcc.</code>		
Arguments	VARIABLES : <code>collection(var-dvar)</code> VALUES : <code>collection(val-int, omax-int)</code>		
Restrictions	<code>required(VARIABLES, var)</code> <code> VALUES > 0</code> <code>required(VALUES, [val, omax])</code> <code>increasing_seq(VALUES, [val])</code> <code>VALUES.ymax ≥ 0</code> <code>VALUES.ymax ≤ VARIABLES </code>		
Purpose	For each $i \in [1, VALUES]$, the values of the corresponding set of values $VALUES[j].val$ ($i \leq j \leq VALUES $) should be taken by at most $VALUES[i].ymax$ variables of the <code>VARIABLES</code> collection. From that previous definition, the <code>ymax</code> attributes are decreasing.		
Example	$\left(\begin{array}{l} \langle 2, 0, 1, 0, 0 \rangle, \\ \langle val - 0 \ ymax - 5, val - 1 \ ymax - 3, val - 2 \ ymax - 1 \rangle \end{array} \right)$		
	The <code>ordered_global_cardinality</code> constraint holds since the values of the three sets of values $\{0, 1, 2\}$, $\{1, 2\}$ and $\{2\}$ are respectively used no more than 5, 3 and 1 times within the collection $\langle 2, 0, 1, 0, 0 \rangle$.		
Symmetry	Items of <code>VARIABLES</code> are permutable .		
Arg. properties	Contractible wrt. <code>VALUES</code> .		
Usage	The <code>ordered_global_cardinality</code> can be used in order to restrict the way we assign the values of the <code>VALUES</code> collection to the variables of the <code>VARIABLES</code> collection. It expresses the fact that, when we use a value v , we implicitly also use all values that are less than or equal to v . As depicted by Figure 5.628 this is for instance the case for a <i>soft cumulative</i> constraint where we want to control the shape of cumulative profile by providing for each instant i a variable h_i that gives the height of the cumulative profile at instant i . These variables h_i are passed as the first argument of the <code>ordered_global_cardinality</code> constraint. Then the <code>ymax</code> attribute of the j -th item of the <code>VALUES</code> collection gives the maximum number of instants for which the height of the cumulative profile is greater than or equal to value $VALUES[j].val$. In Figure 5.628 we should have:		

- no more than 1 height variable greater than or equal to 2,
- no more than 3 height variables greater than or equal to 1,
- no more than 5 height variables greater than or equal to 0.




Figure 5.628: (A) A cumulative profile wrt two tasks ① and ②, and its corresponding height variables h_1, h_2, \dots, h_5 giving at each instant how many resource is used (B) profile of value utilisation of the height variables (e.g., value 1 is assigned to variables h_3, h_2, h_4 and therefore used three times)

Remark

The original definition of the `ordered_global_cardinality` constraint mentions a third argument, namely the minimum number of occurrences of the smallest value. We omit it since it is redundant.

An other closely related constraint, the `cost_ordered_global_cardinality` constraint was introduced in [312] in order to model the fact that overloads costs may depend of the instant where they occur.

Algorithm

A filtering algorithm achieving `arc-consistency` in $O(|VARIABLES| + |VALUES|)$ is described in [312]. It is based on the equivalence between the following two statements:

1. the `ordered_global_cardinality` constraint has a solution,
2. all variables of the `VARIABLES` collection assigned to their respective minimum value correspond to a solution to the `ordered_global_cardinality` constraint.

Reformulation

The `ordered_global_cardinality`($\langle var - V_1, var - V_2, \dots, var - V_{|VARIABLES|} \rangle$, $\langle val - v_1 \text{ omax} - o_1, val - v_2 \text{ omax} - o_2, \dots, val - v_{|VALUES|} \text{ omax} - o_{|VALUES|} \rangle$) constraint can be reformulated into a `global_cardinality`($\langle var - V_1, var - V_2, \dots, var - V_{|VARIABLES|} \rangle$, $\langle val - v_1 \text{ nooccurrence} - N_1, val - v_2 \text{ nooccurrence} - N_2, \dots, val - v_{|VALUES|} \text{ nooccurrence} - N_{|VALUES|} \rangle$) and $|VALUES|$ sliding linear inequalities constraints of the form:

$$\begin{aligned}
 N_1 + N_2 + \dots + N_{|VALUES|} &\leq o_1, \\
 N_2 + \dots + N_{|VALUES|} &\leq o_2, \\
 &\dots, \\
 N_{|VALUES|} &\leq o_{|VALUES|}.
 \end{aligned}$$

However, with the next example, T. Petit and J.-C. Régin have shown that this reformulation hinders propagation:

1. $V_1 \in \{0, 1\}, V_2 \in \{0, 1\}, V_3 \in \{0, 1, 2\}, V_4 \in \{2, 3\}, V_5 \in \{2, 3\}$.
2. `global_cardinality`($\langle V_1, V_2, V_3, V_4, V_5 \rangle, \langle \text{val} - 1 \text{ nooccurrence} - N_1, \text{val} - 2 \text{ nooccurrence} - N_2, \text{val} - 3 \text{ nooccurrence} - N_3 \rangle$),
3. $N_1 + N_2 + N_3 \leq 3 \wedge N_2 + N_3 \leq 2 \wedge N_3 \leq 2$.

The previous reformulation does not remove value 2 from the domain of variable V_3 .

See also

related: `cumulative` (*controlling the shape of the cumulative profile for breaking symmetry*), `global_cardinality_low-up`, `increasing_global_cardinality` (*the order is imposed on the main variables, and not on the count variables*).

root concept: `global_cardinality`.

Keywords

application area: assignment.

constraint type: value constraint, order constraint.

filtering: arc-consistency.

Arc input(s)

For all items of VALUES:

VARIABLES

Arc generator $SELF \mapsto \text{collection}(\text{variables})$ **Arc arity**

1

Arc constraint(s) $\text{variables.var} \geq \text{VALUES.val}$ **Graph property(ies)** $\mathbf{NVERTEX} \leq \text{VALUES.ymax}$ **Graph model**

Since we want to express one unary constraint for each value we use the “For all items of VALUES” iterator. Part (A) of Figure 5.629 shows the initial graphs associated with each value 0, 1 and 2 of the VALUES collection of the **Example** slot. Part (B) of Figure 5.629 shows the corresponding final graph associated with value 0. Since we use the **NVERTEX** graph property, the vertices of the final graph is stressed in bold.




Figure 5.629: Initial and final graph of the ordered_global_cardinality constraint