

5.179 `in_interval_reified`

	DESCRIPTION	LINKS
Origin	Reified version of <code>in_interval</code> .	
Constraint	<code>in_interval_reified(VAR, LOW, UP, B)</code>	
Synonyms	<code>dom_reified</code> , <code>in_reified</code> .	
Arguments	VAR : <code>dvar</code> LOW : <code>int</code> UP : <code>int</code> B : <code>dvar</code>	
Restrictions	$LOW \leq UP$ $B \geq 0$ $B \leq 1$	
Purpose	Enforce the following equivalence, $VAR \in [LOW, UP] \Leftrightarrow B$.	
Example	<div style="border: 1px solid black; padding: 2px; display: inline-block;">(3, 2, 5, 1)</div> The <code>in_interval_reified</code> constraint holds since: <ul style="list-style-type: none"> • Its first argument $VAR = 3$ is greater than or equal to its second argument $LOW = 2$ and less than or equal to its third argument $UP = 5$ (i.e., $3 \in [2, 5]$). • The corresponding Boolean variable B is set to 1 since condition $3 \in [2, 5]$ holds. 	
Typical	$VAR \neq LOW$ $VAR \neq UP$ $LOW < UP$	
Symmetries	<ul style="list-style-type: none"> • An occurrence of a value of VAR that belongs to $[LOW, UP]$ (resp. does not belong to $[LOW, UP]$) can be replaced by any other value in $[LOW, UP]$ (resp. not in $[LOW, UP]$). • One and the same constant can be added to VAR, LOW and UP. 	
Reformulation	The <code>in_interval_reified</code> constraint can be reformulated in terms of linear constraints. For convenience, we rename VAR to x , LOW to l , UP to u , and B to y . The constraint is decomposed into the following conjunction of constraints: $x \geq l \Leftrightarrow y_1,$ $x \leq u \Leftrightarrow y_2,$ $y_1 \wedge y_2 \Leftrightarrow y.$	

$$x \geq l \Leftrightarrow y_1,$$

$$x \leq u \Leftrightarrow y_2,$$

$$y_1 \wedge y_2 \Leftrightarrow y.$$

We show how to encode these constraints with linear inequalities. The first constraint, i.e., $x \geq l \Leftrightarrow y_1$ is encoded by posting one of the following three constraints:

$$\begin{cases} \text{a)} & \text{if } \underline{x} \geq l : y_1 = 1, \\ \text{b)} & \text{if } \bar{x} < l : y_1 = 0, \\ \text{c)} & \text{otherwise : } x \geq (l - \underline{x}) \cdot y_1 + \underline{x} \wedge x \leq (\bar{x} - l + 1) \cdot y_1 + l - 1. \end{cases}$$

On the one hand, cases a) and b) correspond to situations where one can fix y_1 , no matter what value will be assigned to x . On the other hand, in case c), y_1 can take both values 0 or 1 depending on the value assigned to x . As shown by Figure 5.404, all possible solutions for the pair of variables (x, y_1) satisfy the following two linear inequalities $x \geq (l - \underline{x}) \cdot y_1 + \underline{x}$ and $x \leq (\bar{x} - l + 1) \cdot y_1 + l - 1$. The first inequality discards all points that are above the line that goes through the two extreme solution points $(\underline{x}, 0)$ and $(l, 1)$, while the second one removes all points that are below the line that goes through the two extreme solution points $(l - 1, 0)$ and $(\bar{x}, 1)$.




Figure 5.404: Illustration of the reformulation of the reified constraint $x \geq l \Leftrightarrow y_1$ with two linear inequalities

The second constraint, i.e., $x \leq u \Leftrightarrow y_2$ is encoded by posting one of the following three constraints:

$$\begin{cases} \text{d)} & \text{if } \bar{x} \leq u : y_2 = 1, \\ \text{e)} & \text{if } \underline{x} > u : y_2 = 0, \\ \text{f)} & \text{otherwise : } x \leq (u - \bar{x}) \cdot y_2 + \bar{x} \wedge x \geq (\underline{x} - u - 1) \cdot y_2 + u + 1. \end{cases}$$

On the one hand, cases d) and e) correspond to situations where one can fix y_2 , no matter what value will be assigned to x . On the other hand, in case f), y_2 can take both value 0 or 1 depending on the value assigned to x . As shown by Figure 5.405, all possible solutions for the pair of variables (x, y_2) satisfy the following two linear inequalities $x \leq (u - \bar{x}) \cdot y_2 + \bar{x}$ and $x \geq (\underline{x} - u - 1) \cdot y_2 + u + 1$. The first inequality discards all points that are above the line that goes through the two extreme solution points $(\bar{x}, 0)$ and $(u, 1)$, while the second one removes all points that are below the line that goes through the two extreme solution points $(u + 1, 0)$ and $(\underline{x}, 1)$.




Figure 5.405: Illustration of the reformulation of the reified constraint $x \leq u \Leftrightarrow y_2$ with two linear inequalities

The third constraint, i.e., $y_1 \wedge y_2 \Leftrightarrow y$ is encoded as:

$$\begin{cases} \text{g)} & y \geq y_1 + y_2 - 1, \\ \text{h)} & y \leq y_1, \\ \text{i)} & y \leq y_2. \end{cases}$$

Case g) handles the implication $y_1 \wedge y_2 \Rightarrow y$, while cases h) and i) take care of the other side $y \Rightarrow y_1 \wedge y_2$.

See also

specialisation: `in_interval`.

uses in its reformulation: `alldifferent` (*bound consistency preserving reformulation*).

Keywords

characteristic of a constraint: reified constraint.

constraint arguments: binary constraint.

constraint type: predefined constraint, value constraint.

filtering: arc-consistency.

20100916

1341